



Two generalizations of some fixed point theorems

Ovidiu Popescu

Department of Mathematical Analysis and Probability, Transilvania University, Iuliu Maniu 50, RO-505801, Brasov, Romania

ARTICLE INFO

Article history:

Received 7 June 2011

Received in revised form 19 September 2011

Accepted 19 September 2011

Keywords:

Fixed point

Contraction

Nonexpansive map

Greguš's theorem

ABSTRACT

The aim of this paper is to generalize two classical fixed point theorems given by Bogin [J. Bogin, A generalization of a fixed point theorem of Goebel, Kirk and Shimi, Canad. Math. Bull. 19 (1976) 7–12] and Greguš [M. Greguš, A fixed point theorem in Banach spaces, Boll. Un. Math. Ital. A (5) 17 (1980) 193–198]. We also complement and extend some very recent results proved by Suzuki [T. Suzuki, A generalized Banach contraction principle that characterizes metric completeness, Proc. Amer. Math. Soc. 136 (2008) 1861–1869].

© 2011 Elsevier Ltd. All rights reserved.

1. Introduction

Let (X, d) be a metric space, T a self-mapping on X and k a nonnegative real number such that the inequality

$$d(Tx, Ty) \leq kd(x, y)$$

holds for any $x, y \in X$. If $k < 1$ then T is said to be a contractive mapping, if $k = 1$, then T is said to be a nonexpansive mapping. The well-known Banach theorem states that if X is complete, then every contractive mapping has a unique fixed point and that point can be obtained as a limit of repeated iteration of the mapping at any point of X . However, a nonexpansive mapping need not have fixed points. Yet these mappings have a fixed point when X has a convex structure. There exists a huge literature about contractive and nonexpansive type mappings, where the contractive and nonexpansive conditions are replaced with more general conditions (see, for instance [1–13]).

Bogin [1] proved the following result.

Theorem 1.1. *Let (X, d) be a nonempty complete metric space and $T : X \rightarrow X$ a mapping satisfying*

$$d(Tx, Ty) \leq ad(x, y) + b[d(x, Tx) + d(y, Ty)] + c[d(x, Ty) + d(y, Tx)], \quad (1)$$

where $a \geq 0$, $b > 0$, $c > 0$ and $a + 2b + 2c = 1$. Then T has a unique fixed point.

This result was generalized by Ćirić [3] and Li [14]. Greguš [9] considered a class of self-mapping T on X which satisfy (1) and (2) with $c = 0$. He proved the following theorem.

Theorem 1.2. *Let C be a nonempty closed convex subset of a Banach space B and $T : C \rightarrow C$ a mapping that satisfies*

$$\|Tx - Ty\| \leq a\|x - y\| + b(\|x - Tx\| + \|y - Ty\|) \quad (2)$$

for all $x, y \in C$, where $a > 0$, $b > 0$ and $a + 2b = 1$. Then T has a unique fixed point.

E-mail address: ovidiu.popescu@unitbv.ro.

Greguš's result has inspired many authors in further investigations: Abdeljawad and Karapinar [15], Ćirić [16–18], Delbosco et al. [19], Diviccaro et al. [7], Fisher [8], Fisher and Sessa [20], Jungck [10], Li [14], Murthy et al. [21], Mukherjee and Verma [22], Raswan and Ahmed [23], Ćirić [24] has constructed an example to show that if the mapping T satisfies (1) with $b = 0$ and if a and c are such that (2) holds, then T need not have a fixed point.

The following remarkable generalization of the classical Banach contraction theorem, due to Suzuki [25], has lead to some important contributions in metric fixed point theory (see, for instance, [26–33]).

Theorem 1.3. Let (X, d) be a complete metric space and $S : X \rightarrow X$. Define a nonincreasing function θ from $[0, 1)$ onto $(\frac{1}{2}, 1]$ by

$$\theta(r) = \begin{cases} 1 & \text{if } 0 \leq r \leq (\sqrt{5} - 1)/2, \\ (1 - r)/r^2 & \text{if } (\sqrt{5} - 1)/2 \leq r \leq 1/\sqrt{2}, \\ 1/(1 + r) & \text{if } 1/\sqrt{2} \leq r < 1. \end{cases}$$

Assume that there exists $r \in [0, 1)$ such that $\theta(r)d(x, Sx) \leq d(x, y)$ implies $d(Sx, Sy) \leq rd(x, y)$, for all $x, y \in X$. Then S has a unique fixed point.

Also, Kikkawa and Suzuki [34] proved Kannan and Meir–Keeler's versions of Theorem 1.3. Moreover, Suzuki studied a class of operators satisfying the following condition.

Definition 1.4. Let T be a mapping on a subset C of a Banach space E . Then T is said to satisfy “condition” (C) if for all $x, y \in C$

$$(C) \quad 1/2 \|x - Tx\| \leq \|x - y\| \text{ implies } \|Tx - Ty\| \leq \|x - y\|.$$

The condition (C) is weaker than nonexpansiveness.

Inspired by these results we give in this paper two generalizations of Theorems 1.1 and 1.2.

2. Main results

The first result of this paper is the following generalization of Bogin's theorem.

Theorem 2.1. Let (X, d) be a nonempty complete metric space and $T : X \rightarrow X$ be a mapping satisfying

$$1/2d(x, Tx) \leq d(x, y)$$

implies

$$d(Tx, Ty) \leq ad(x, y) + b[d(x, Tx) + d(y, Ty)] + c[d(x, Ty) + d(y, Tx)], \quad (3)$$

where $a \geq 0$, $b > 0$, $c > 0$ and $a + 2b + 2c = 1$. Then T has a unique fixed point.

Proof. Let $x \in X$ be arbitrary. Since $1/2d(x, Tx) \leq d(x, Tx)$ we have that

$$d(Tx, T^2x) \leq ad(x, Tx) + b[d(x, Tx) + d(Tx, T^2x)] + c[d(x, T^2x) + d(Tx, Tx)].$$

Hence

$$d(Tx, T^2x) \leq (a + b)d(x, Tx) + bd(Tx, T^2x) + c[d(x, Tx) + d(Tx, T^2x)].$$

Therefore we obtain

$$d(Tx, T^2x) \leq \frac{a + b + c}{1 - b - c}d(x, Tx) = d(x, Tx).$$

This implies that the sequence $\{d_n\}_{n=0}^{\infty}$ is a decreasing one, where $d_n := d(T^n x, T^{n+1} x)$ and $T^0 x := x$.

Next we will show that there exists a nonnegative number $m < 2$ such that $d(Tx, T^3x) \leq md_0$. First, we suppose that $d(x, T^2x) \geq d(x, Tx)$. Then $1/2d(x, Tx) \leq d(x, T^2x)$ and we have

$$d(Tx, T^3x) \leq ad(x, T^2x) + bd(x, Tx) + bd(T^2x, T^3x) + cd(x, T^3x) + cd(Tx, T^2x).$$

Thus

$$d(Tx, T^3x) \leq a(d_0 + d_1) + bd_0 + bd_2 + c[d_0 + d(Tx, T^3x)] + cd_1$$

and so,

$$d(Tx, T^3x) \leq \frac{2a + 2b + 2c}{1 - c}d_0 = \frac{1 + a}{1 - c}d_0.$$

Setting $m_1 = (1 + a)/(1 - c)$, we have $m_1 < 2$ and $d(Tx, T^3x) \leq m_1 d_0$. Now, we suppose that $d(x, T^2x) < d(x, Tx)$. Since

$$d(Tx, T^2x) \leq ad(x, Tx) + b[d(x, Tx) + d(Tx, T^2x)] + c[d(x, T^2x) + d(Tx, Tx)],$$

we get

$$d_1 < (a + b)d_0 + bd_1 + cd_0.$$

Hence

$$d_1 < \frac{a + b + c}{1 - b} d_0$$

and then

$$d(Tx, T^3x) \leq d(Tx, T^2x) + d(T^2x, T^3x) \leq 2d(Tx, T^2x) \leq \frac{2a + 2b + 2c}{1 - b} d_0 = \frac{1 + a}{1 - b} d_0.$$

Setting $m_2 = (1 + a)/(1 - b)$, we have $m_2 < 2$ and $d(Tx, T^3x) \leq m_2 d_0$. Taking $m = \max\{m_1, m_2\}$, we get $0 < m < 2$ and $d(Tx, T^3x) \leq md(x, Tx)$, for all $x \in X$.

Since $1/2d(Tx, T^2x) \leq d(Tx, T^2x)$ we have

$$d(T^2x, T^3x) \leq ad(Tx, T^2x) + b[d(Tx, T^2x) + d(T^2x, T^3x)] + cd(Tx, T^3x).$$

Thus

$$d_2 \leq (a + 2b)d_0 + mcd_0 = (a + 2b + mc)d_0.$$

Setting $k = a + 2b + mc$, we have $k < 1$ and $d_2 \leq kd_0$ for all $x \in X$.

Let $x_0 \in X$ and $u_n = T^n x_0$. Then $d_{n+2} \leq kd_n$ for all $n \geq 0$, where $d_n = d(u_n, u_{n+1})$. Therefore, for any even integer $n \geq 0$ we have by induction $d_n \leq k^{n/2} d_0 \leq k^{(n-1)/2} d_0$ and for every odd integer $n \geq 1$ we have also by induction $d_n \leq k^{(n-1)/2} d_1 \leq k^{(n-1)/2} d_0$. Hence, for all $n \geq 0$ we get $d_n \leq k^{(n-1)/2} d_0$. Since $k \in (0, 1)$ we obtain that $\{u_n\}$ is a Cauchy sequence and by completeness of X we have that there exists $z \in X$ such that the sequence $\{u_n\}$ converges to z as $n \rightarrow \infty$.

Next, we will show that z is a fixed point of T . Assuming that there exists n such that $d(z, u_n) < 1/2d(u_n, u_{n+1})$ and $d(z, u_{n+1}) < 1/2d(u_{n+1}, u_{n+2})$ we obtain

$$d_n = d(u_n, u_{n+1}) \leq d(z, u_n) + d(z, u_{n+1}) < 1/2(d_n + d_{n+1}) \leq d_n.$$

This is a contradiction, so for all $n \geq 0$ we have either $d(z, u_n) \geq 1/2d(u_n, u_{n+1})$ or $d(z, u_{n+1}) \geq 1/2d(u_{n+1}, u_{n+2})$. Thus, there exists a subsequence $\{n_j\}$ of $\{n\}$ such that $d(u_{n_j}, z) \geq 1/2d(u_{n_j+1}, u_{n_j})$ for every integer $j \geq 0$. Then, we have

$$d(Tz, u_{n_j+1}) \leq ad(z, u_{n_j}) + bd(z, Tz) + bd(u_{n_j}, u_{n_j+1}) + cd(z, u_{n_j+1}) + cd(Tz, u_{n_j}).$$

Taking $j \rightarrow \infty$ we get $d(Tz, z) \leq (b + c)d(Tz, z)$. This implies $d(Tz, z) = 0$ and so, $Tz = z$.

If y is another fixed point T then $d(y, z) \geq 1/2d(z, Tz) = 0$ and then

$$d(y, z) = d(Ty, Tz) \leq ad(y, z) + b[d(y, y) + d(z, z)] + c[d(y, z) + d(y, z)].$$

Hence $d(y, z) \leq (a + 2c)d(y, z)$. This implies $d(y, z) = 0$, which is a contradiction. So, T has a unique fixed point. \square

Example 2.2. Let $X = [-1, 1]$ with the usual metric and let $T : X \rightarrow X$ be given as

$$Tx = \begin{cases} -x & \text{if } x \in [0, 3/4) \cup (3/4, 1] = U, \\ x/2 & \text{if } x \in [-1, 0) = V, \\ 0 & \text{if } x = 3/4. \end{cases}$$

We will prove that:

1. T has a unique fixed point.
2. T satisfies condition (3) with $a = 1/2$, $b = c = 1/8$, i.e.,

$$d(x, Tx)/2 \leq d(x, y) \Rightarrow d(Tx, Ty) \leq M(x, y),$$

where $M(x, y) = \frac{1}{2}d(x, y) + \frac{1}{8}[d(x, Tx) + d(y, Ty) + d(x, Ty) + d(y, Tx)]$.

3. T does not satisfy Suzuki's condition from Theorem 1.3.

4. T does not satisfy Bogin's condition (1) with $a = 1/2$, $b = c = 1/8$.

Proof. 1 is obvious. 2.

- (i) For $x, y \in U$ it is $M(x, y) = \frac{1}{2}|y - x| + \frac{1}{8}(2x + 2y + x + y + x + y) = \frac{1}{2}|y - x| + \frac{1}{2}(x + y) \geq |y - x| = d(Tx, Ty)$ and (3) holds.
- (ii) If $x, y \in V$, then $M(x, y) \geq \frac{1}{2}d(x, y) = \frac{1}{2}|y - x| = d(Tx, Ty)$, so (3) holds.
- (iii) If $x \in U, y \in V$, then $M(x, y) = \frac{1}{2}(x - y) + \frac{1}{8}(2x - \frac{1}{2}y + x - \frac{1}{2}y + |y + x|) = \frac{7}{8}x - \frac{5}{8}y + \frac{1}{8}|y + x| \geq x - \frac{1}{2}y, d(Tx, Ty) = |x + \frac{1}{2}y|$. Since $x \geq 0, y < 0$ we have $x - \frac{1}{2}y \geq x + \frac{1}{2}y$ and $x - \frac{1}{2}y \geq -x - \frac{1}{2}y$, so (3) holds.
- (iv) If $x \in V, y \in U$, then $M(x, y) \geq d(Tx, Ty)$ like in (iii).
- (v) For $x \in U, y = \frac{3}{4}$, it is $M(x, y) = \frac{1}{2}|x - \frac{3}{4}| + \frac{1}{8}(2x + \frac{3}{4} + x + \frac{3}{4} + x) = \frac{1}{2}|x - \frac{3}{4}| + \frac{1}{2}x + \frac{3}{16}$ and $d(Tx, Ty) = x$. Since $d(x, Tx)/2 \leq d(x, y)$ we have $x \leq |x - \frac{3}{4}|$, so $x \leq \frac{3}{8}$. Therefore $M(x, y) = \frac{3}{8} + \frac{3}{16} = \frac{9}{16} \geq x$, and (3) holds.
- (vi) For $x \in V, y = \frac{3}{4}$, it is $M(x, y) = \frac{1}{2}(\frac{3}{4} - x) + \frac{1}{8}(-\frac{x}{2} + \frac{3}{4} - x + \frac{3}{4} - \frac{x}{2}) = -\frac{3x}{4} + \frac{9}{16} \geq \frac{1}{2}$, and $d(Tx, Ty) = -\frac{x}{2} \leq \frac{1}{2}$. Then (3) holds.
- (vii) If $x = \frac{3}{4}, y \in U$, then $M(x, y) = \frac{1}{2}|\frac{3}{4} - y| + \frac{1}{8}(\frac{3}{4} + 2y + y + \frac{3}{4} + y) = \frac{1}{2}|\frac{3}{4} - y| + \frac{y}{2} + \frac{3}{16}$ and $d(Tx, Ty) = y$. By $d(x, Tx)/2 \leq d(x, y)$ we have $\frac{3}{8} \leq |\frac{3}{4} - y|$, so $y \leq \frac{3}{8}$. Therefore $M(x, y) = \frac{3}{8} + \frac{3}{16} = \frac{9}{16} \geq y$. Hence (3) holds.
- (viii) If $x = \frac{3}{4}, y \in V$, then $M(x, y) = \frac{1}{2}(\frac{3}{4} - y) + \frac{1}{8}(\frac{3}{4} - \frac{y}{2} - y + \frac{3}{4} - \frac{y}{2}) = \frac{9}{16} - \frac{3y}{4} \geq \frac{1}{2}$ and $d(Tx, Ty) = -\frac{y}{2} \leq \frac{1}{2}(\frac{y}{2} \in [-\frac{1}{2}, 0])$. Hence (3) holds.
- (ix) If $x = y$ then (3) is obvious.

3. If $x = 0, y = 1$, then $\theta(r)d(x, Tx) = 0 < 1 = d(x, y)$ and $d(Tx, Ty) = 1$, so condition from Theorem 1.3. does not hold.

4. If $x = \frac{3}{4}, y = 1$ we have $d(Tx, Ty) = 1$ and $M(x, y) = \frac{1}{8} + \frac{1}{8}(\frac{3}{4} + 2 + 1 + 1 + \frac{3}{4}) = \frac{13}{16}$, so $d(Tx, Ty) > M(x, y)$. Therefore Bogin's condition (2) does not hold. \square

For the next theorem we need recall an important concept.

Definition 2.3 (Takahashi [35]). Let (X, d) be a metric space. A mapping $W : X \times X \times [0, 1] \rightarrow X$ is said to be a *convex structure* on X if for each $(x, y, \lambda) \in X \times X \times [0, 1]$ and $u \in X$

$$d(u, W(x, y, \lambda)) \leq \lambda d(u, x) + (1 - \lambda)d(u, y).$$

The metric space X together with W is called a *convex metric space*.

It is obvious that in a convex metric space we have $d(x, W(x, y, \lambda)) = (1 - \lambda)d(x, y)$ and $d(y, W(x, y, \lambda)) = \lambda d(x, y)$. Now we can prove a generalization of Gregus' s theorem.

Theorem 2.4. Let X be a complete convex metric space and let $T : X \rightarrow X$ be a selfmapping. Assume that $1/2d(x, Tx) \leq d(x, y)$ implies

$$d(Tx, Ty) \leq ad(x, y) + bd(x, Tx) + bd(y, Ty), \quad (4)$$

where $a > 0, b > 0$ and $a + 2b = 1$. Then T has a unique fixed point.

Proof. Let $x_0 \in X, u_n = T^n x_0$ and $d_n = d(u_n, u_{n+1})$. Since $1/2d(T^n x_0, T^{n+1} x_0) \leq d(T^n x_0, T^{n+1} x_0)$, we have from the assumption that

$$d_{n+1} \leq ad_n + bd_n + bd_{n+1}.$$

Thus

$$d_{n+1} \leq \frac{a+b}{1-b}d_n = d_n,$$

for all $n \geq 0$. So, $\{d_n\}$ is a decreasing sequence.

Let $t = W(Tx_0, T^2x_0, 1/2)$ and $z = W(T^2x_0, T^3x_0, 1/2)$. Suppose that $d(Tx_0, z) < 1/2d(Tx_0, T^2x_0)$. Since

$$d(T^2x_0, z) = 1/2d(T^2x_0, T^3x_0) \leq 1/2d(Tx_0, T^2x_0)$$

it follows that

$$d(Tx_0, T^2x_0) \leq d(Tx_0, z) + d(T^2x_0, z) < d(Tx_0, T^2x_0),$$

which is a contradiction. Thus we have $d(Tx_0, z) \geq 1/2d(Tx_0, T^2x_0)$. From the assumption we obtain

$$d(T^2x_0, Tz) \leq ad(Tx_0, z) + bd(Tx_0, T^2x_0) + bd(z, Tz).$$

Since $1/2d(T^2x_0, T^3x_0) = d(T^2x_0, z) \leq d(T^2x_0, z)$, we also obtain

$$d(T^3x_0, Tz) \leq ad(T^2x_0, z) + bd(T^2x_0, T^3x_0) + bd(z, Tz).$$

Therefore

$$\begin{aligned} d(z, Tz) &\leq 1/2d(Tz, T^2x_0) + 1/2d(Tz, T^3x_0) \\ &\leq 1/2[ad(Tx_0, z) + bd(Tx_0, T^2x_0) + bd(z, Tz)] + 1/2[ad(T^2x_0, z) + bd(T^2x_0, T^3x_0) + bd(z, Tz)]. \end{aligned}$$

Thus we get

$$(2 - 2b)d(z, Tz) \leq ad(z, Tx_0) + ad(z, T^2x_0) + 2bd(x_0, Tx_0),$$

and so,

$$\begin{aligned} d(z, Tz) &\leq \frac{a}{1+a}d(z, Tx_0) + \frac{a}{2(1+a)}d(T^2x_0, T^3x_0) + \frac{2b}{1+a}d_0 \\ &\leq \frac{a}{1+a}d(z, Tx_0) + \frac{a}{2(1+a)}d_0 + \frac{2b}{1+a}d_0 \\ &= \frac{a}{1+a}d(z, Tx_0) + \frac{2-a}{2(1+a)}d_0. \end{aligned}$$

Next we consider two cases.

Case 1. Suppose that $1/2d(x_0, Tx_0) \leq d(x_0, T^2x_0)$. By the assumption we get

$$\begin{aligned} d(T^3x_0, Tx_0) &\leq ad(x_0, T^2x_0) + bd(x_0, Tx_0) + bd(T^2x_0, T^3x_0) \\ &\leq a[d(x_0, Tx_0) + d(Tx_0, T^2x_0)] + 2bd_0 \\ &\leq (2a + 2b)d_0 = (1 + a)d_0. \end{aligned}$$

Since $d(z, Tx_0) \leq 1/2d(Tx_0, T^2x_0) + 1/2d(Tx_0, T^3x_0)$ we obtain $d(z, Tx_0) \leq 1/2(2 + a)d_0$. Therefore

$$d(z, Tz) \leq \frac{a}{1+a} \cdot \frac{2+a}{2}d_0 + \frac{2-a}{2(1+a)}d_0 = \frac{a^2 + a + 2}{2(1+a)}d_0.$$

Since $a^2 + a + 2 < 2 + 2a$, taking

$$k_1 = \frac{a^2 + a + 2}{2(1+a)}$$

we have $k_1 < 1$ and $d(z, Tz) \leq k_1d_0$.

Case 2. Suppose that $1/2d(x_0, Tx_0) > d(x_0, T^2x_0)$. We will show that there exists $k_2 < 1$ such that $d(t, Tt) \leq k_2d_0$. Assume that $d(x_0, t) < 1/2d(x_0, Tx_0)$. If $d(Tx_0, t) \leq 1/2d(x_0, Tx_0)$ then we have $d(x_0, Tx_0) \leq d(x_0, t) + d(Tx_0, t) < d(x_0, Tx_0)$, which is a contradiction. So, we get $d(Tx_0, t) > 1/2d(x_0, Tx_0)$. Since $d(Tx_0, t) = 1/2d(Tx_0, T^2x_0)$, we obtain that $d(Tx_0, T^2x_0) > d(x_0, Tx_0)$, which is also a contradiction. Therefore, we must have $1/2d(x_0, Tx_0) \leq d(x_0, t)$. By the assumption we get

$$d(Tx_0, Tt) \leq ad(x_0, t) + bd(t, Tt) + bd(x_0, Tx_0).$$

Since $1/2d(Tx_0, T^2x_0) = 1/2d(x_0, t) \leq d(x_0, t)$ we also get

$$d(T^2x_0, Tt) \leq ad(Tx_0, t) + bd(t, Tt) + bd(Tx_0, T^2x_0).$$

Hence we obtain

$$\begin{aligned} d(t, Tt) &\leq 1/2d(Tx_0, Tt) + 1/2d(T^2x_0, Tt) \\ &\leq 1/2[ad(x_0, t) + bd(t, Tt) + bd(x_0, Tx_0)] + 1/2[ad(Tx_0, t) + bd(t, Tt) + bd(Tx_0, T^2x_0)]. \end{aligned}$$

Thus

$$\begin{aligned} (1 - b)d(t, Tt) &\leq a/2d(x_0, t) + a/2d(Tx_0, t) + bd(x_0, Tx_0) \\ &= a/2d(x_0, t) + a/4d(Tx_0, T^2x_0) + bd(x_0, Tx_0) \\ &\leq a/2d(x_0, t) + (a/4 + b)d_0. \end{aligned}$$

Since $d(x_0, t) \leq 1/2d(x_0, Tx_0) + 1/2d(x_0, T^2x_0) < 1/2d_0 + 1/4d_0 = 3/4d_0$, we obtain

$$(1 - b)d(t, Tt) \leq (3a/8 + a/4 + b)d_0,$$

which implies

$$d(t, Tt) \leq \frac{5/8a + b}{1 - b} d_0.$$

But $5/8a + b < 1 - b$, so taking

$$k_2 = \frac{5/8a + b}{1 - b}$$

we have $k_2 < 1$ and $d(t, Tt) \leq k_2 d_0$.

Hence in all cases we proved that there exists $k < 1$ and $y \in X$ such that $d(y, Ty) < kd(x_0, Tx_0)$. Therefore $\inf\{d(x, Tx) : x \in X\} = 0$.

Next, we will prove that the infimum is attained. Take the following system of sets:

$K_n := \{x \in X : d(x, Tx) \leq r/(2n)\}$, TK_n and $\overline{TK_n}$, where $\overline{TK_n}$ is the closure of TK_n , $r = (1 - a)/(1 + b)$, $n \geq 1$.

For any $x, y \in K_n$ such that $1/2d(x, Tx) \leq d(x, y)$ we have

$$\begin{aligned} d(x, y) &\leq d(x, Tx) + d(Tx, Ty) + d(y, Ty) \\ &\leq \frac{r}{2n} + ad(x, y) + bd(x, Tx) + bd(y, Ty) + \frac{r}{2n} \\ &\leq ad(x, y) + \frac{(1 + b)r}{n}. \end{aligned}$$

Thus

$$d(x, y) \leq \frac{1 + b}{1 - a} \cdot \frac{r}{n} = \frac{1}{n}.$$

If $1/2d(x, Tx) > d(x, y)$ we have

$$d(x, y) < \frac{1}{2} \cdot \frac{r}{2n} < \frac{1}{n}.$$

Hence $d(x, y) \leq 1/n$ for all $x, y \in K_n$, i.e. $\text{diam}(K_n) \leq 1/n$. Since $d(Tx, T^2x) \leq d(x, Tx)$ we have $T(K_n) \subseteq K_n$ and then, $\text{diam}(\overline{T(K_n)}) = \text{diam}(T(K_n)) \leq 1/n$.

Supposing $y \in \overline{T(K_n)}$, then for any $\epsilon > 0$ there exists $y' \in K_n$ such that $d(y, Ty') < \epsilon$. If $d(y, y') < 1/2d(y', Ty')$ and $d(y, Ty') < 1/2d(Ty', T^2y')$, then

$$d(y', Ty') < d(y, y') + d(y, Ty') < 1/2[d(y', Ty') + d(Ty', T^2y')] \leq d(y', Ty'),$$

which is a contradiction. Hence, either $d(y, y') \geq 1/2d(y', Ty')$ or $d(y, Ty') \geq 1/2d(Ty', T^2y')$. In the first case we have by the assumption

$$d(Ty, Ty') \leq ad(y, y') + bd(y, Ty) + bd(y', Ty').$$

Thus

$$\begin{aligned} d(y, Ty) &\leq d(y, Ty') + d(Ty, Ty') \\ &\leq \epsilon + ad(y, y') + bd(y, Ty) + \frac{br}{2n} \\ &\leq \epsilon + a[d(y, Ty') + d(Ty', y')] + bd(y, Ty) + \frac{br}{2n}. \end{aligned}$$

Hence

$$d(y, Ty) \leq \frac{(a + 1)\epsilon}{1 - b} + \frac{a + b}{1 - b} \cdot \frac{r}{2n} < \frac{a + 3}{1 - b}\epsilon + \frac{r}{2n}.$$

In the second case, from the assumption we obtain

$$d(Ty, T^2y') \leq ad(y, Ty') + bd(y, Ty) + bd(Ty', T^2y').$$

But $d(Ty', T^2y') \leq 2d(y, Ty') = 2\epsilon$ and then

$$\begin{aligned} d(y, Ty) &\leq d(y, T^2y') + d(T^2y', Ty) \leq d(y, Ty') + d(Ty', T^2y') + d(T^2y', Ty) \\ &\leq \epsilon + 2\epsilon + ad(y, Ty') + bd(y, Ty) + \frac{br}{2n} \\ &\leq \epsilon + 2\epsilon + a\epsilon + bd(y, Ty) + \frac{br}{2n} \\ &\leq (a+3)\epsilon + bd(y, Ty) + \frac{br}{2n}. \end{aligned}$$

Hence

$$d(y, Ty) \leq \frac{a+3}{1-b}\epsilon + \frac{b}{1-b} \cdot \frac{r}{2n} < \frac{a+3}{1-b}\epsilon + \frac{r}{2n}.$$

Therefore, in all cases we proved that

$$d(y, Ty) \leq \frac{a+3}{1-b}\epsilon + \frac{r}{2n}.$$

Since $\epsilon > 0$ is arbitrary, we get that $d(y, Ty) \leq r/(2n)$, i.e. $y \in K_n$. Hence we have $\overline{T(K_n)} \subseteq K_n$. Therefore, $\{\overline{T(K_n)}\}$ is a decreasing sequence of closed nonempty sets with $\text{diam}(\overline{T(K_n)}) \rightarrow 0$ as $n \rightarrow \infty$. Thus they have a nonempty intersection. Since $\overline{T(K_n)} \subseteq K_n$, we obtain that there exists $u \in K_n$ for all n . This implies that u is a fixed point of T .

If v is another fixed point of T , since $1/2d(u, Tu) = 0 \leq d(u, v)$ we have

$$d(u, v) = d(Tu, Tv) \leq ad(u, v) + bd(u, u) + bd(v, v) = ad(u, v),$$

which is a contradiction. \square

Example 2.5. Let $X = [-1, 1]$ with the usual metric and let $T : X \rightarrow X$ be given as

$$Tx = \begin{cases} -x & \text{if } x \in [0, 2/3) \cup (2/3, 1] = U, \\ x/2 & \text{if } x \in [-1, 0) = V, \\ -1/4 & \text{if } x = 2/3. \end{cases}$$

We will prove that:

1. T has a unique fixed point.
2. T satisfies condition (4) with $a = 2/3$, $b = 1/6$, i.e.,

$$d(x, Tx)/2 \leq d(x, y) \Rightarrow d(Tx, Ty) \leq m(x, y),$$
 where $m(x, y) = \frac{2}{3}d(x, y) + \frac{1}{6}[d(x, Tx) + d(y, Ty)]$.
3. T does not satisfy Suzuki's condition from Theorem 1.3.
4. T does not satisfy Greguš's condition (2) with $a = 2/3$, $b = 1/6$.

Proof. 1 is obvious. 2.

- (i) For $x, y \in U$ it is $m(x, y) = \frac{2}{3}|y - x| + \frac{2x}{6} + \frac{2y}{6} \geq \frac{2}{3}|y - x| + \frac{1}{3}|y - x| = |y - x| = d(Tx, Ty)$ and (4) holds.
- (ii) If $x, y \in V$, then $m(x, y) = \frac{2}{3}|y - x| + \frac{1}{6}|\frac{x}{2}| + \frac{1}{6}|\frac{y}{2}| \geq \frac{1}{2}|y - x| = d(Tx, Ty)$, so (4) holds.
- (iii) If $x \in U$, $y \in V$, then $m(x, y) = \frac{2}{3}|y - x| + \frac{2x}{6} + \frac{1}{6}|\frac{y}{2}| = \frac{2}{3}(x - y) + \frac{1}{3}x - \frac{1}{12}y = x - \frac{3}{4}y$, $d(Tx, Ty) = |x + \frac{1}{2}y|$. Since $x \geq 0, y < 0$ we have $x - \frac{3}{4}y \geq x + \frac{1}{2}y$ and $x - \frac{3}{4}y \geq -x - \frac{1}{2}y$, so (4) holds.
- (iv) If $x \in V$, $y \in U$, then $m(x, y) \geq d(Tx, Ty)$ like in (iii).
- (v) For $x \in U$, $y = \frac{2}{3}$, it is $m(x, y) = \frac{2}{3}|x - \frac{2}{3}| + \frac{2x}{6} + \frac{1}{6}(\frac{2}{3} + \frac{1}{4})$ and $d(Tx, Ty) = |x - \frac{1}{4}|$. Since $d(x, Tx)/2 \leq d(x, y)$ we have $x \leq |x - \frac{2}{3}|$, so $x \leq \frac{1}{3}$. Therefore $|x - \frac{1}{4}| \leq \frac{1}{4}$ and $|x - \frac{2}{3}| \geq \frac{1}{3}$. Hence $m(x, y) \geq \frac{2}{9} + \frac{1}{6}(\frac{2}{3} + \frac{1}{4}) = \frac{3}{8} > d(Tx, Ty)$, and (4) holds.
- (vi) For $x \in V$, $y = \frac{2}{3}$, it is $m(x, y) = \frac{2}{3}(\frac{2}{3} - x) + \frac{1}{6}|\frac{x}{2}| + \frac{1}{6}(\frac{2}{3} + \frac{1}{4}) \geq \frac{4}{9} > \frac{1}{4}$, and $d(Tx, Ty) = |\frac{x}{2} + \frac{1}{4}| \leq \frac{1}{4}$ ($\frac{x}{2} \in [-\frac{1}{2}, 0)$), so (4) holds.
- (vii) If $x = \frac{2}{3}$, $y \in U$, then $m(x, y) = \frac{2}{3}|\frac{2}{3} - y| + \frac{1}{6}(\frac{2}{3} + \frac{1}{4}) + \frac{2y}{6}$ and $d(Tx, Ty) = |y - \frac{1}{4}|$. By $d(x, Tx)/2 \leq d(x, y)$ we have $\frac{1}{2}(\frac{2}{3} + \frac{1}{4}) = \frac{11}{24} \leq |y - \frac{2}{3}|$, so $y \leq \frac{5}{24}$. Therefore $m(x, y) \geq \frac{2}{3} \times \frac{11}{24} > \frac{1}{4}$ and $d(Tx, Ty) \leq \frac{1}{4}$. Hence (4) holds.
- (viii) If $x = \frac{2}{3}$, $y \in V$, then $m(x, y) \geq \frac{2}{3}d(x, y) = \frac{2}{3}(\frac{2}{3} - y) \geq \frac{4}{9} > \frac{1}{4}$ and $d(Tx, Ty) = |-\frac{1}{4} - \frac{y}{2}| \leq \frac{1}{4}$ ($\frac{y}{2} \in [-\frac{1}{2}, 0)$). Hence (4) holds.
- (ix) If $x = y$ then (4) is obvious.

3. If $x = 0, y = 1$, then $\theta(r)d(x, Tx) = 0 < 1 = d(x, y)$ and $d(Tx, Ty) = 1$, so condition from Theorem 1.3. does not hold.

4. If $x = \frac{2}{3}, y = 1$ we have $d(Tx, Ty) = \frac{3}{4}$ and $m(x, y) = \frac{2}{9} + \frac{2}{6} + \frac{1}{6}(\frac{2}{3} + \frac{1}{4}) = \frac{51}{72} < \frac{54}{72} = \frac{3}{4}$, so $d(Tx, Ty) > m(x, y)$. Therefore Greguš's condition (2) does not hold. \square

References

- [1] J. Bogin, A generalization of a fixed point theorem of Goebel, Kirk and Shimi, *Canad. Math. Bull.* 19 (1976) 7–12.
- [2] S.K. Chatterjea, Applications of an extension of a theorem of Disher on common fixed point, *Pure Math. Manuscript* 6 (1987) 35–38.
- [3] Lj.B. Ćirić, On some discontinuous fixed point mappings in convex metric spaces, *Czechoslovak Math. J.* 43 (118) (1993) 319–326.
- [4] Lj.B. Ćirić, Nonexpansive type mappings and fixed point theorems in convex metric spaces, *Rend. Accad. Naz. Sci. XI Mem. Mat. Appl.* (5) XIX (1995) 263–271.
- [5] Lj.B. Ćirić, On some nonexpansive type mappings and fixed point, *Indian J. Pure Appl. Math.* 24 (3) (1993) 145–149.
- [6] Lj.B. Ćirić, On some mappings in metric space and fixed points, *Acad. Roy. Belg. Bull. Cl. Sci.* (5) (1995) 81–89.
- [7] M.L. Diviccaro, B. Fisher, S. Sessa, A common fixed point theorem of Greguš type, *Publ. Math. Debrecen* 34 (1997) 83–89.
- [8] B. Fisher, Common fixed points on a Banach space, *Chung Yuan J.* 11 (1982) 19–26.
- [9] M. Greguš, A fixed point theorem in Banach spaces, *Boll. Unione Mat. Ital. Sez. A* (5) 17 (1980) 193–198.
- [10] G. Jungck, On a fixed point theorem of Fisher and Sessa, *Int. J. Math. Math. Sci.* 13 (1990) 497–500.
- [11] R. Kannan, Some results on fixed point theory-II, *Amer. Math. Monthly* 76 (1969) 405–408.
- [12] E. Karapinar, K. Tas, Generalized (C)-conditions and related fixed point theorems, *Comput. Math. Appl.* 61 (11) (2011) 3370–3380.
- [13] T. Zamfirescu, Fixed point theorems in metric spaces, *Arch. Math.* 23 (1972) 292–298.
- [14] B.I. Li, Fixed point theorems of nonexpansive mappings in convex metric spaces, *Appl. Math. Mech.* 10 (1989) 183–188.
- [15] T. Abdeljawad, E. Karapinar, A common fixed point theorem of a Greguš type on convex cone metric spaces, *J. Comput. Anal. Appl.* 13 (4) (2011) 609–621.
- [16] Lj.B. Ćirić, On a common fixed point theorem of a Greguš type, *Publ. Inst. Math.* 49 (1991) 174–178.
- [17] Lj.B. Ćirić, On Diviccaro, Fisher and Sessa open questions, *Arch. Math.* 29 (1993) 145–152.
- [18] Lj.B. Ćirić, On a generalization of a Greguš fixed point theorem, *Czechoslovak Math. J.* 50 (2000) 449–458.
- [19] D. Delbosco, O. Ferrero, F. Rossati, Teoremi di punto fisso per applicazioni negli spazi di Banach, *Boll. Un. Mat. Ital. Sez. A* (6) 2 (1993) 297–303.
- [20] B. Fisher, S. Sessa, On a fixed point theorem of Greguš, *Int. J. Math. Math. Sci.* 9 (1986) 23–28.
- [21] P.P. Murty, Y.J. Cho, B. Fisher, Common fixed points of Greguš type mappings, *Glas. Mat.* 30 (50) (1995) 335–341.
- [22] R.N. Mukherjee, V. Verma, A note on a fixed point of Greguš, *Math. Japon.* 33 (1988) 745–749.
- [23] R.A. Rashwan, M.A. Ahmed, Common fixed points of Greguš type multi-valued mappings, *Arch. Math.* 38 (2002) 37–47.
- [24] Lj.B. Ćirić, A new class of nonexpansive type mappings and fixed points, *Czechoslovak Math. J.* 49 (124) (1999) 891–899.
- [25] T. Suzuki, A generalized Banach contraction principle that characterizes metric completeness, *Proc. Amer. Math. Soc.* 136 (2008) 1861–1869.
- [26] S. Dhompangsa, H. Yingtaeweessittikul, Fixed points for multivalued mappings and the metric completeness, *Fixed Point Theory Appl.* (2009) 1–15. Article ID 972395.
- [27] Gh. Mot, A. Petrusel, Fixed point theory for a new type of contractive multivalued operators, *Nonlinear Anal.* 70 (2009) 3371–3377.
- [28] O. Popescu, Two fixed point theorems for generalized contractions with constants in complete metric space, *Cent. Eur. J. Math.* 7 (3) (2009) 529–538.
- [29] M. Kikkawa, T. Suzuki, Some similarity between contractions and Kannan mappings, *Fixed Point Theory Appl.* (2008) 1–8. Article ID 649749.
- [30] S.L. Singh, S.N. Mishra, Coincidence theorems for certain classes of hybrid contractions, *Fixed Point Theory Appl.* (2010) 1–14. Article ID 898109.
- [31] T. Suzuki, Some remarks on recent generalization of the Banach contraction principle, in: *Proc. of the 8th Int. Conf. on Fixed Point Theory and its Appl.*, 2007, pp. 751–761.
- [32] T. Suzuki, Fixed point theorems and convergence theorems for some generalized nonexpansive mappings, *J. Math. Anal. Appl.* 340 (2008) 1088–1095.
- [33] T. Suzuki, A new type of fixed point theorem in metric spaces, *Nonlinear Anal.* 71 (2009) 5313–5317.
- [34] M. Kikkawa, T. Suzuki, Three fixed point theorems for generalized contractions with constants in complete metric spaces, *Nonlinear Anal.* 69 (2008) 2942–2949.
- [35] W. Takahashi, A convexity in metric space and nonexpansive mappings, *Kodai Math. Sem. Rep.* 22 (1970) 142–149.